

NUP43 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57576

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	Q8NFH3
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42151
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NUP43
Epitope Specificity	1-100/380
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus; nuclear pore complex.
SIMILARITY	Contains 6 WD repeats.
SUBUNIT	Component of the Nup107-160 subcomplex of the nuclear pore complex (NPC). The Nup107-160 subcomplex includes NUP160, NUP133, NUP107, NUP98, NUP85, NUP43, NUP37, SEH1 and SEC13.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Bidirectional transport of macromolecules between the cytoplasm and nucleus occurs through nuclear pore complexes (NPCs) embedded in the nuclear envelope. NPCs are composed of subcomplexes, and NUP43 is part of one such subcomplex, Nup107-160 (Loiodice et al., 2004 [PubMed 15146057]).[supplied by OMIM, Mar 2008]


Additional Information

Gene ID	348995
Other Names	Nucleoporin Nup43, Nup107-160 subcomplex subunit Nup43, p42, NUP43
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	NUP43
Function	Component of the Nup107-160 subcomplex of the nuclear pore complex (NPC). The Nup107-160 subcomplex is required for the assembly of a functional NPC. The Nup107-160 subcomplex is also required for normal kinetochore microtubule attachment, mitotic progression and chromosome segregation.
Cellular Location	Chromosome, centromere, kinetochore. Nucleus, nuclear pore complex

Images



Sample:
Hela(Human) Cell Lysate at 30 ug
293T(Human) Cell Lysate at 30 ug
Primary: Anti-NUP43 (AP57576) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 42 kD
Observed band size: 30 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NUP43) Polyclonal Antibody, Unconjugated (AP57576) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.