

PTPN22 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57589

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9Y2R2
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	91705

Additional Information

Gene ID	26191
Other Names	Tyrosine-protein phosphatase non-receptor type 22, 3.1.3.48, Hematopoietic cell protein-tyrosine phosphatase 70Z-PEP, Lymphoid phosphatase, LyP, PEST-domain phosphatase, PEP, PTPN22, PTPN8
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	PTPN22
Synonyms	PTPN8
Function	Acts as a negative regulator of T-cell receptor (TCR) signaling by direct dephosphorylation of the Src family kinases LCK and FYN, ITAMs of the TCRz/CD3 complex, as well as ZAP70, VAV, VCP and other key signaling molecules (PubMed: 16461343 , PubMed: 18056643). Associates with and probably dephosphorylates CBL. Dephosphorylates LCK at its activating 'Tyr-394' residue (PubMed: 21719704). Dephosphorylates ZAP70 at its activating 'Tyr-493' residue (PubMed: 16461343). Dephosphorylates the immune system activator SKAP2 (PubMed: 21719704). Positively regulates toll-like receptor (TLR)-induced type 1 interferon production (PubMed: 23871208). Promotes host antiviral responses mediated by type 1 interferon (By similarity). Regulates NOD2-induced pro-inflammatory cytokine secretion and autophagy (PubMed: 23991106). Acts as an activator of NLRP3

inflammasome assembly by mediating dephosphorylation of 'Tyr-861' of NLRP3 (PubMed:[27043286](#)). Dephosphorylates phospho-anandamide (p-AEA), an endocannabinoid to anandamide (also called N-arachidonylethanolamide) (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P29352}.

Tissue Location

Expressed in bone marrow, B and T-cells, PBMCs, natural killer cells, monocytes, dendritic cells and neutrophils (PubMed:15208781). Both isoform 1 and 4 are predominantly expressed in lymphoid tissues and cells. Isoform 1 is expressed in thymocytes and both mature B and T-cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.