

SF20 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57617

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q969H8
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18795
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MYDGF
Epitope Specificity	51-150/173
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted. Endoplasmic reticulum-Golgi intermediate compartment.
SIMILARITY	Belongs to the UPF0556 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene was previously thought to support proliferation of lymphoid cells and was considered an interleukin. However, this activity has not been reproducible and the function of this protein is currently unknown. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	56005
Other Names	Myeloid-derived growth factor {ECO:0000303 PubMed:25581518, ECO:0000312 HGNC:HGNC:16948}, MYDGF, MYDGF (HGNC:16948)
Target/Specificity	Expressed in synovial tissue. Found in synovial fluid of patients with arthropaties.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MYDGF (HGNC:16948)
Function	Bone marrow-derived monocyte and paracrine-acting protein that promotes cardiac myocyte survival and adaptive angiogenesis for cardiac protection and/or repair after myocardial infarction (MI). Stimulates endothelial cell proliferation through a MAPK1/3-, STAT3- and CCND1-mediated signaling pathway. Inhibits cardiac myocyte apoptosis in a PI3K/AKT-dependent signaling pathway (By similarity). Involved in endothelial cell proliferation and angiogenesis (PubMed: 25581518).
Cellular Location	Secreted. Endoplasmic reticulum-Golgi intermediate compartment. Endoplasmic reticulum. Golgi apparatus. Note=The C-terminal RTEL motif may provide retention in the endoplasmic reticulum
Tissue Location	Expressed in eosinophils (at protein level) (PubMed:29954947). Expressed in bone marrow cells (PubMed:25581518) Expressed in synovial tissue. Found in synovial fluid of patients with arthropaties (PubMed:17362502).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.