

SF3A3 Rabbit pAb

SF3A3 Rabbit pAb
Catalog # AP57619

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q12874
Predicted	Human, Mouse, Rat, Dog, Pig, Horse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58849
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SF3A3
Epitope Specificity	21-120/501
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus speckle.
SIMILARITY	Belongs to the SF3A3 family.
SUBUNIT	Identified in the spliceosome C complex. Component of splicing factor SF3A which is composed of three subunits; SF3A3/SAP61, SF3A2/SAP62, SF3A1/SAP114. SF3A associates with the splicing factor SF3B and a 12S RNA unit to form the U2 small nuclear ribonucleoproteins complex (U2 snRNP). Interacts with SF3A1, through its N-terminal region.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes subunit 3 of the splicing factor 3a protein complex. The splicing factor 3a heterotrimer includes subunits 1, 2 and 3 and is necessary for the in vitro conversion of 15S U2 snRNP into an active 17S particle that performs pre-mRNA splicing. Subunit 3 interacts with subunit 1 through its amino-terminus while the zinc finger domain of subunit 3 plays a role in its binding to the 15S U2 snRNP. This gene has a pseudogene on chromosome 20. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	10946
Other Names	Splicing factor 3A subunit 3, SF3a60, Spliceosome-associated protein 61, SAP 61, SF3A3, SAP61
Target/Specificity	Ubiquitous.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SF3A3
Synonyms	SAP61
Function	Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed: 10882114 , PubMed: 11533230 , PubMed: 32494006 , PubMed: 34822310 , PubMed: 8022796). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed: 10882114 , PubMed: 11533230 , PubMed: 32494006 , PubMed: 34822310). Within the 17S U2 SnRNP complex, SF3A3 is part of the SF3A subcomplex that contributes to the assembly of the 17S U2 snRNP, and the subsequent assembly of the pre-spliceosome 'E' complex and the pre-catalytic spliceosome 'A' complex (PubMed: 10882114 , PubMed: 11533230). Involved in pre-mRNA splicing as a component of pre-catalytic spliceosome 'B' complexes (PubMed: 29360106 , PubMed: 30315277).
Cellular Location	Nucleus speckle. Nucleus
Tissue Location	Ubiquitous..

Background

This gene encodes subunit 3 of the splicing factor 3a protein complex. The splicing factor 3a heterotrimer includes subunits 1, 2 and 3 and is necessary for the in vitro conversion of 15S U2 snRNP into an active 17S particle that performs pre-mRNA splicing. Subunit 3 interacts with subunit 1 through its amino-terminus while the zinc finger domain of subunit 3 plays a role in its binding to the 15S U2 snRNP. This gene has a pseudogene on chromosome 20. [provided by RefSeq, Jul 2008]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.