10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

QSOX2 Rabbit pAb

QSOX2 Rabbit pAb
Catalog # AP57627

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q6ZRP7
Reactivity	Mouse
Predicted	Rat, Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	77529
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human QSOX2
Epitope Specificity	101-200/698
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane. Secreted. Cell membrane. Nucleus membrane. Seems to be predominantly targeted to the nuclear and outer plasma membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	QSOX2 is a member of the sulfhydryl oxidase/quiescin-6 (Q6) family (QSOX1; MIM 603120) that regulates the sensitization of neuroblastoma cells for IFN-gamma (IFNG; MIM 147570)-induced cell death (Wittke et al., 2003 [PubMed 14633699]).[supplied by OMIM, Jun 2009]

Additional Information

Gene ID	169714
Other Names	Sulfhydryl oxidase 2, 1.8.3.2, Neuroblastoma-derived sulfhydryl oxidase, Quiescin Q6-like protein 1, QSOX2, QSCN6L1, SOXN
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	QSOX2 (HGNC:30249)

Synonyms	QSCN6L1, SOXN
Function	Sulfhydryl oxidase that catalyzes the oxidation of protein thiol groups to form disulfide bonds, with the reduction of oxygen to hydrogen peroxide (By similarity). May contribute to disulfide bond formation in secreted proteins (By similarity). May play a role in regulating the sensitization of neuroblastoma cells for interferon- gamma-induced apoptosis (PubMed:14633699). Required for normal ovarian function (By similarity).
Cellular Location	Membrane; Single- pass membrane protein. Secreted. Cell membrane; Single-pass membrane protein. Nucleus membrane; Single- pass membrane protein. Note=Seems to be predominantly targeted to the nuclear and outer plasma membrane
Tissue Location	Expressed in pancreas, brain, placenta, kidney, heart and fetal tissues. Weakly expressed in lung, liver and skeletal muscles.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (QSOX2) Polyclonal Antibody, Unconjugated (AP57627) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.