

RAD51AP1 Rabbit pAb

RAD51AP1 Rabbit pAb

Catalog # AP57640

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q96B01
Predicted	Rat, Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38457
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RAD51AP1
Epitope Specificity	1-100/352
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Colocalizes with RAD51 to multiple nuclear foci.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	10635
Other Names	RAD51-associated protein 1, HsRAD51AP1, RAD51-interacting protein, RAD51AP1 {ECO:0000303 PubMed:16990250, ECO:0000312 HGNC:HGNC:16956}
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	RAD51AP1 {ECO:0000303 PubMed:16990250, ECO:0000312 HGNC:HGNC:16956}
Function	Structure-specific DNA-binding protein involved in DNA repair by promoting RAD51-mediated homologous recombination (PubMed: 17996710 , PubMed: 17996711 , PubMed: 20871616 , PubMed: 25288561 , PubMed: 26323318). Acts by stimulating D-Loop formation by RAD51:

specifically enhances joint molecule formation through its structure-specific DNA interaction and its interaction with RAD51 (PubMed:[17996710](#), PubMed:[17996711](#)). Binds single-stranded DNA (ssDNA), double-stranded DNA (dsDNA) and secondary DNA structures, such as D-loop structures: has a strong preference for branched-DNA structures that are obligatory intermediates during joint molecule formation (PubMed:[17996710](#), PubMed:[17996711](#), PubMed:[22375013](#), PubMed:[9396801](#)). Cooperates with WDR48/UAF1 to stimulate RAD51-mediated homologous recombination: both WDR48/UAF1 and RAD51AP1 have coordinated role in DNA-binding during homologous recombination and DNA repair (PubMed:[27239033](#), PubMed:[27463890](#), PubMed:[32350107](#)). WDR48/UAF1 and RAD51AP1 also have a coordinated role in DNA-binding to promote USP1-mediated deubiquitination of FANCD2 (PubMed:[31253762](#)). Also involved in meiosis by promoting DMC1-mediated homologous meiotic recombination (PubMed:[21307306](#)). Key mediator of alternative lengthening of telomeres (ALT) pathway, a homology-directed repair mechanism of telomere elongation that controls proliferation in aggressive cancers, by stimulating homologous recombination (PubMed:[31400850](#)). May also bind RNA; additional evidences are however required to confirm RNA-binding in vivo (PubMed:[9396801](#)).

Cellular Location

Chromosome. Nucleus Chromosome, telomere. Note=Colocalizes with RAD51 to multiple nuclear foci (By similarity). Colocalizes with DMC1 on meiotic chromatin (By similarity) {ECO:0000250|UniProtKB:Q8C551}

Tissue Location

Highly expressed in testis and thymus (PubMed:9396801). Lower levels in colon and small intestine (PubMed:9396801). Little or no expression in spleen, prostate, ovary and peripheral blood leukocytes (PubMed:9396801)

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.