

PCBP2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57710

Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q15366</u>

Reactivity Rat, Pig, Bovine

HostRabbitClonalityPolyclonalCalculated MW38580Physical StateLiquid

Immunogen KLH conjugated synthetic peptide derived from human PCBP2

Epitope Specificity 1-100/365 **Isotype** IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus. Cytoplasm. Loosely bound in the nucleus. May shuttle between the

nucleus and the cytoplasm.

SIMILARITY Contains 3 KH domains.

SUBUNIT Identified in a mRNP complex, at least composed of DHX9,DDX3X, ELAVL1,

HNRNPU, IGF2BP1, ILF3, PABPC1, PCBP2, PTBP2, STAU1, STAU2, SYNCRIP and YBX1. Interacts with IFIH1 and RNF135. Interacts with MAVS (via C-terminus)

and ITCH (via WW domains).

Post-translational Phosphorylated. The non-phosphorylated form(s) exhibited the strongest

modifications poly(rC)-binding activity.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions The protein encoded by this gene appears to be multifunctional. Along with

PCBP-1 and hnRNPK, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been

replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Jul 2008]

Additional Information

Gene ID 5094

Other Names Poly(rC)-binding protein 2, Alpha-CP2, Heterogeneous nuclear

ribonucleoprotein E2, hnRNP E2, PCBP2

Target/Specificity Detected in all tissues examined.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name PCBP2 {ECO:0000303 | PubMed:7607214, ECO:0000312 | HGNC:HGNC:8648}

Function Single-stranded nucleic acid binding protein that binds preferentially to oligo

dC (PubMed:<u>12414943</u>, PubMed:<u>7607214</u>). Major cellular poly(rC)-binding protein (PubMed:<u>12414943</u>). Also binds poly(rU) (PubMed:<u>12414943</u>). Acts as

a negative regulator of antiviral signaling (PubMed: 19881509,

PubMed:<u>35322803</u>). Negatively regulates cellular antiviral responses mediated by MAVS signaling (PubMed:<u>19881509</u>). It acts as an adapter between MAVS and the E3 ubiquitin ligase ITCH, therefore triggering MAVS ubiquitination and degradation (PubMed:<u>19881509</u>). Negativeley regulates the cGAS-STING pathway via interaction with CGAS, preventing the formation of liquid- like droplets in which CGAS is activated (PubMed:<u>35322803</u>).

Together with PCBP1, required for erythropoiesis, possibly by regulating


mRNA splicing (By similarity).

Cellular Location Nucleus. Cytoplasm. Note=Loosely bound in the nucleus (PubMed:7607214).

May shuttle between the nucleus and the cytoplasm (PubMed:7607214).

Tissue Location Detected in all tissues examined.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCBP2) Polyclonal Antibody, Unconjugated (AP57710) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.