

PCQAP Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57711

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q96RN5
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86753

Additional Information

Gene ID	51586
Other Names	Mediator of RNA polymerase II transcription subunit 15, Activator-recruited cofactor 105 kDa component, ARC105, CTG repeat protein 7a, Mediator complex subunit 15, Positive cofactor 2 glutamine/Q-rich-associated protein, PC2 glutamine/Q-rich-associated protein, TPA-inducible gene 1 protein, TIG-1, Trinucleotide repeat-containing gene 7 protein, MED15, ARC105, CTG7A, PCQAP, TIG1, TNRC7
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MED15
Synonyms	ARC105, CTG7A, PCQAP, TIG1, TNRC7
Function	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene- specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for cholesterol- dependent gene regulation. Positively regulates the Nodal signaling pathway.
Cellular Location	Cytoplasm. Nucleus.

Tissue Location

Expressed in all tissues examined, including heart, brain, lung, spleen, thymus, pancreas, blood leukocyte and placenta However, the level of expression varied, with highest expression in the placenta and peripheral blood and lowest in the pancreas and kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.