

TMEM109 Rabbit pAb

TMEM109 Rabbit pAb

Catalog # AP57727

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9BVC6
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26210
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TMEM109
Epitope Specificity	151-243/243
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus outer membrane. Endoplasmic reticulum membrane. Sarcoplasmic reticulum membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	79073
Other Names	Voltage-gated monoatomic cation channel TMEM109, TMEM109 (HGNC:28771)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	TMEM109 (HGNC:28771)
Function	Functions as a voltage-gated monoatomic cation channel permeable to both potassium and calcium (By similarity). Plays a role in the cellular response to DNA damage (PubMed: 23542032).
Cellular Location	Nucleus outer membrane {ECO:0000250 UniProtKB:O77751}; Multi-pass

membrane protein {ECO:0000250|UniProtKB:O77751}. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:O77751}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O77751}. Sarcoplasmic reticulum membrane {ECO:0000250|UniProtKB:O77751}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O77751}

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:

U251(Human) Cell Lysate at 40 ug


Panc-1(Human) Cell Lysate at 40 ug

Primary: Anti-TMEM109 (AP57727) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 23 kD

Observed band size: 23 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TMEM109) Polyclonal Antibody, Unconjugated (AP57727) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.