

PPP1R11 Rabbit pAb

PPP1R11 Rabbit pAb

Catalog # AP57744

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	O15091
Reactivity	Rat
Predicted	Pig, Dog, Human, Mouse, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67315
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PPP1R11
Epitope Specificity	21-100/126
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 14 contains about 700 genes and 106 million base pairs and makes up about 3.5% of human cellular DNA. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease. The SERPINA1 gene is located on chromosome 14 and when defective leads to the genetic disorder α -antitrypsin deficiency. This disorder is characterized by severe lung complications and liver dysfunction. Notably, the immunoglobulin heavy chain locus is found on chromosome 14 and has been identified as a fusion with the chromosome 19 encoded protein BCL3 in the (14;19) translocations found in a variety of B cell malignancies. The KIAA0391 gene product has been provisionally designated KIAA0391 pending further characterization.

Additional Information

Gene ID	9692
Other Names	Mitochondrial ribonuclease P catalytic subunit, 3.1.26.5, Mitochondrial ribonuclease P protein 3, Mitochondrial RNase P protein 3, Protein only RNase P catalytic subunit {ECO:0000312 HGNC:HGNC:19958}, PRORP (HGNC:19958)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	PRORP (HGNC:19958)
Function	Catalytic ribonuclease component of mitochondrial ribonuclease P, a complex composed of TRMT10C/MRPP1, HSD17B10/MRPP2 and PRORP/MRPP3, which cleaves tRNA molecules in their 5'-ends (PubMed: 18984158 , PubMed: 25953853 , PubMed: 34715011). The presence of TRMT10C/MRPP1, HSD17B10/MRPP2 is required to catalyze tRNA molecules in their 5'-ends (PubMed: 25953853).
Cellular Location	Mitochondrion.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat stomach tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PPP1R11) Polyclonal Antibody, Unconjugated (AP57744) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.