

RanBP3 Rabbit pAb

RanBP3 Rabbit pAb
Catalog # AP57765

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9H6Z4
Reactivity	Rat
Predicted	Human, Mouse, Chicken, Dog, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60210
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RanBP3
Epitope Specificity	441-540/567
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
SIMILARITY	Contains 1 RanBD1 domain.
SUBUNIT	Interacts with CHC1 in a Ran-stimulated manner. Interacts with XPO1. Interacts (via its C-terminal R domain) with SMAD2 (dephosphorylated form via its MH1 and MH2 domains); the interaction results in the nuclear export of SMAD2 and termination of the TGF-beta signaling. Interacts (via its C-terminal R domain) with SMAD3 (dephosphorylated form via its MH1 domain); the interaction results in the nuclear export of SMAD3 and termination of the TGF-beta signaling.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Acts as a cofactor for XPO1/CRM1-mediated nuclear export, perhaps as export complex scaffolding protein. Bound to XPO1/CRM1, stabilizes the XPO1/CRM1-cargo interaction. In the absence of Ran-bound GTP prevents binding of XPO1/CRM1 to the nuclear pore complex. Binds to CHC1/RCC1 and increases the guanine nucleotide exchange activity of CHC1/RCC1. Recruits XPO1/CRM1 to CHC1/RCC1 in a Ran-dependent manner. Negative regulator of TGF-beta signaling through interaction with the R-SMAD proteins, SMAD2 and SMAD3, and mediating their nuclear export.

Additional Information

Gene ID	8498
Other Names	Ran-binding protein 3, RanBP3, RANBP3
Target/Specificity	Widely expressed with high levels in testis and heart.

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	RANBP3
Function	Acts as a cofactor for XPO1/CRM1-mediated nuclear export, perhaps as export complex scaffolding protein. Bound to XPO1/CRM1, stabilizes the XPO1/CRM1-cargo interaction. In the absence of Ran-bound GTP prevents binding of XPO1/CRM1 to the nuclear pore complex. Binds to CHC1/RCC1 and increases the guanine nucleotide exchange activity of CHC1/RCC1. Recruits XPO1/CRM1 to CHC1/RCC1 in a Ran-dependent manner. Negative regulator of TGF-beta signaling through interaction with the R-SMAD proteins, SMAD2 and SMAD3, and mediating their nuclear export.
Cellular Location	Cytoplasm. Nucleus. Note=Nuclear import is promoted by phosphorylation at Ser-126 and is dependent on KPNA4.
Tissue Location	Widely expressed with high levels in testis and heart.

Background

Acts as a cofactor for XPO1/CRM1-mediated nuclear export, perhaps as export complex scaffolding protein. Bound to XPO1/CRM1, stabilizes the XPO1/CRM1-cargo interaction. In the absence of Ran-bound GTP prevents binding of XPO1/CRM1 to the nuclear pore complex. Binds to CHC1/RCC1 and increases the guanine nucleotide exchange activity of CHC1/RCC1. Recruits XPO1/CRM1 to CHC1/RCC1 in a Ran-dependent manner. Negative regulator of TGF-beta signaling through interaction with the R-SMAD proteins, SMAD2 and SMAD3, and mediating their nuclear export.

Images



Tissue/cell: Rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
 Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
 Incubation: Anti-RanBP3 Polyclonal Antibody, Unconjugated(AP57765) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.