

COUP-TF1 Rabbit pAb

COUP-TF1 Rabbit pAb

Catalog # AP57796

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P10589
Reactivity	Rat
Predicted	Pig, Human, Mouse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46156
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human COUP-TF1
Epitope Specificity	21-120/423
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	COUP transcription factors have been cloned in several species and identified as orphan members of the steroid/thyroid hormone receptor superfamily. COUP-TFI (also designated COUP or EAR-3) and ARP-1 (also designated COUP-TFII) exhibit highly regulated and overlapping expression in most tissues. COUP-TFs are highly expressed in the developing and central nervous system, suggesting that these factors may be important in neural development and differentiation. COUP-TFs can compete for binding to response elements which are common to other members of this family, including RAR, RXR, PPAR, HNF-4, VDR and TR. They have been shown to act as negative regulators as well as initiators of transcription.

Additional Information

Gene ID	7025
Other Names	COUP transcription factor 1, COUP-TF1, COUP transcription factor I, COUP-TF I, Nuclear receptor subfamily 2 group F member 1, V-erbA-related protein 3, EAR-3, NR2F1, EAR3, ERBAL3, TFCOUP1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	NR2F1
Synonyms	EAR3, ERBAL3, TFCOUP1
Function	Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. Binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Represses transcriptional activity of LHCG.
Cellular Location	Nucleus.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COUP-TF1) Polyclonal Antibody, Unconjugated (AP57796) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.