

RYBP Rabbit pAb

RYBP Rabbit pAb
Catalog # AP57804

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8N488
Reactivity	Human, Mouse
Predicted	Rat, Chicken, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24822
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human AAP1
Epitope Specificity	31-130/228
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cytoplasm. Nucleus, nucleoplasm. Note=Primarily found in the nucleus. Detected in a punctate pattern likely to represent Polycomb group (PcG) bodies
SIMILARITY	Contains 1 RanBP2-type zinc finger.
SUBUNIT	Monomer. Interacts with apoptin, DEDD, FADD, CASP8, CASP10, YY1 and GABPB1. Component of repressive BCOR complex containing Polycomb group subcomplex at least composed of BCOR, PCGF1, RING1 and RNF2/RING2. Together with GABPB1 and YY1, it forms a ternary complex, probably being the bridge factor between these two transcription factors. Interacts with CBX2, YAF2, RING1 and RNF2. Interacts with ubiquitin and ubiquitinated proteins. Interacts with ubiquitinated histone H2A (By similarity). Interacts with MDM2, and thereby inhibits ubiquitination of TP53. Identified in a ternary complex containing MDM2, TP53 and RYBP.
Post-translational modifications	Monoubiquitinated
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	May be implicated in the regulation of the transcription as a repressor of the transcriptional activity of E4TF1. In tumor cell lines, may induce apoptosis.

Additional Information

Gene ID	23429
Other Names	RING1 and YY1-binding protein, Apoptin-associating protein 1, APAP-1, Death effector domain-associated factor, DED-associated factor, YY1 and E4TF1-associated factor 1, RYBP, DEDAF, YEAF1
Target/Specificity	Widely expressed with highest levels in lymphoid tissues and placenta.

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	RYBP
Synonyms	DEDAF, YEAF1
Function	Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1-like complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed: 25519132). Component of a PRC1-like complex that mediates monoubiquitination of histone H2A 'Lys-119' on the X chromosome and is required for normal silencing of one copy of the X chromosome in XX females. May stimulate ubiquitination of histone H2A 'Lys-119' by recruiting the complex to target sites (By similarity). Inhibits ubiquitination and subsequent degradation of TP53, and thereby plays a role in regulating transcription of TP53 target genes (PubMed: 19098711). May also regulate the ubiquitin-mediated proteasomal degradation of other proteins like FANK1 to regulate apoptosis (PubMed: 14765135 , PubMed: 27060496). May be implicated in the regulation of the transcription as a repressor of the transcriptional activity of E4TF1 (PubMed: 11953439). May bind to DNA (By similarity). May play a role in the repression of tumor growth and metastasis in breast cancer by down-regulating SRRM3 (PubMed: 27748911).
Cellular Location	Nucleus. Cytoplasm. Nucleus, nucleoplasm {ECO:0000250 UniProtKB:Q8CCI5}. Note=Primarily found in the nucleus Detected in a punctate pattern likely to represent Polycomb group (PcG) bodies (By similarity). {ECO:0000250 UniProtKB:Q8CCI5}
Tissue Location	Down-regulated in breast cancer tissues and in several breast cancer cell lines (at protein level) (PubMed: 27748911) Widely expressed with highest levels in lymphoid tissues and placenta

Background

May be implicated in the regulation of the transcription as a repressor of the transcriptional activity of E4TF1. In tumor cell lines, may induce apoptosis.

Images

Paraformaldehyde-fixed, paraffin embedded (mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RYBP) Polyclonal Antibody, Unconjugated (AP57804) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit)



(sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.