

# PPIP5K2/HISPPD1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57850

#### **Product Information**

**Application** IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>043314</u>

**Reactivity** Rat, Pig, Dog, Bovine

HostRabbitClonalityPolyclonalCalculated MW140407

## **Additional Information**

**Gene ID** 23262

Other Names Inositol hexakisphosphate and diphosphoinositol-pentakisphosphate kinase

2, 2.7.4.21, 2.7.4.24, Diphosphoinositol pentakisphosphate kinase 2, Histidine acid phosphatase domain-containing protein 1, InsP6 and PP-IP5 kinase 2, VIP1 homolog 2, hsVIP2, PPIP5K2 {ECO:0000303|PubMed:29590114,

ECO:0000312 | HGNC:HGNC:29035 }

**Dilution** IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Protein Information**

Name PPIP5K2 {ECO:0000303 | PubMed:29590114,

ECO:0000312 | HGNC:HGNC:29035}

**Function** Bifunctional inositol kinase that acts in concert with the IP6K kinases IP6K1,

IP6K2 and IP6K3 to synthesize the diphosphate group-containing inositol pyrophosphates diphosphoinositol pentakisphosphate, PP-InsP5, and bis-diphosphoinositol tetrakisphosphate, (PP)2-InsP4 (PubMed:17690096, PubMed:17702752, PubMed:21222653, PubMed:29590114). PP-InsP5 and (PP)2-InsP4, also respectively called InsP7 and InsP8, regulate a variety of cellular processes, including apoptosis, vesicle trafficking, cytoskeletal dynamics, exocytosis, insulin signaling and neutrophil activation

(PubMed: 17690096, PubMed: 17702752, PubMed: 21222653, PubMed: 29590114) Phosphorylates inosital bayakisphosphate

PubMed: <u>29590114</u>). Phosphorylates inositol hexakisphosphate (InsP6) at position 1 to produce PP-InsP5 which is in turn phosphorylated by IP6Ks to

produce (PP)2-InsP4 (PubMed:<u>17690096</u>, PubMed:<u>17702752</u>). Alternatively, phosphorylates PP-InsP5 at position 1, produced by IP6Ks from InsP6, to produce (PP)2-InsP4 (PubMed:<u>17690096</u>, PubMed:<u>17702752</u>). Required for normal hearing (PubMed:<u>29590114</u>).

#### **Cellular Location**

Cytoplasm, cytosol.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.