

# QRSL Rabbit pAb

QRSL Rabbit pAb  
Catalog # AP57881

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF
<b>Primary Accession</b>	<a href="#">Q9H0R6</a>
<b>Predicted</b>	Human, Mouse, Rat, Dog, Pig, Sheep
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	57460
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human QRSL
<b>Epitope Specificity</b>	401-500/528
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
<b>SUBCELLULAR LOCATION</b>	Mitochondrion.
<b>SIMILARITY</b>	Belongs to the amidase family. GatA subfamily.
<b>SUBUNIT</b>	Subunit of the heterotrimeric GatCAB amidotransferase (AdT) complex, composed of A (QRSL1), B (GATB) and C (GATC) subunits.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	QRSL belongs to the amidase family. Catalytic activity: ATP + L-glutamyl-tRNA(Gln) + L-glutamine = ADP + phosphate + L-glutaminyl-tRNA(Gln) + L-glutamate. There are two named isoforms.

## Additional Information

---

<b>Gene ID</b>	55278
<b>Other Names</b>	Glutamyl-tRNA(Gln) amidotransferase subunit A, mitochondrial {ECO:0000255 HAMAP-Rule:MF_03150}, Glu-AdT subunit A {ECO:0000255 HAMAP-Rule:MF_03150}, 6.3.5.7 {ECO:0000255 HAMAP-Rule:MF_03150}, Glutaminyl-tRNA synthase-like protein 1 {ECO:0000255 HAMAP-Rule:MF_03150}, QRSL1 {ECO:0000255 HAMAP-Rule:MF_03150}
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	QRSL1 {ECO:0000255   HAMAP-Rule:MF_03150}
<b>Function</b>	Allows the formation of correctly charged Gln-tRNA(Gln) through the transamidation of misacylated Glu-tRNA(Gln) in the mitochondria. The reaction takes place in the presence of glutamine and ATP through an activated gamma-phospho-Glu-tRNA(Gln).
<b>Cellular Location</b>	Mitochondrion {ECO:0000255   HAMAP-Rule:MF_03150, ECO:0000269   PubMed:19805282}

## Background

---

QRSL belongs to the amidase family. Catalytic activity: ATP + L-glutamyl-tRNA(Gln) + L-glutamine = ADP + phosphate + L-glutaminyl-tRNA(Gln) + L-glutamate. There are two named isoforms.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.