

SLC25A1 Rabbit pAb

SLC25A1 Rabbit pAb
Catalog # AP57930

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P53007
Reactivity	Rat, Human
Predicted	Pig, Dog, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34013
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SLC25A1
Epitope Specificity	1-100/311
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
SUBCELLULAR LOCATION	Mitochondrion inner membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the mitochondrial carrier subfamily of solute carrier proteins. Members of this family include nuclear-encoded transporters that translocate small metabolites across the mitochondrial membrane. This protein regulates the movement of citrate across the inner membranes of the mitochondria. Mutations in this gene have been associated with combined D-2- and L-2-hydroxyglutaric aciduria. Pseudogenes of this gene have been identified on chromosomes 7, 11, 16, and 19. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013]

Additional Information

Gene ID	6576
Other Names	Tricarboxylate transport protein, mitochondrial, Citrate transport protein, CTP, Mitochondrial citrate carrier, CIC, Solute carrier family 25 member 1, Tricarboxylate carrier protein, SLC25A1, SLC20A3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	SLC25A1
Synonyms	SLC20A3
Function	Mitochondrial electroneutral antiporter that exports citrate from the mitochondria into the cytosol in exchange for malate (PubMed: 26870663 , PubMed: 29031613 , PubMed: 29238895 , PubMed: 39881208 , PubMed: 38937634). Also able to mediate the exchange of citrate for isocitrate, phosphoenolpyruvate, cis-aconitate and to a lesser extent trans-aconitate, maleate and succinate (PubMed: 29031613). Substrate exchange across the membrane occurs consecutively with one substrate being transported first, then dissociating from the substrate binding site before the second substrate binds for transport in the opposite direction (PubMed: 38937634). In the cytoplasm, citrate plays important roles in fatty acid and sterol synthesis, regulation of glycolysis, protein acetylation, and other physiopathological processes (PubMed: 29031613 , PubMed: 29238895 , PubMed: 39881208).
Cellular Location	Mitochondrion inner membrane {ECO:0000250 UniProtKB:Q8JZU2}; Multi-pass membrane protein


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Jurkat(Human) Cell Lysate at 30 ug
Primary: Anti- SLC25A1 (AP57930) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 34 kD
Observed band size: 34 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLC25A1) Polyclonal Antibody, Unconjugated (AP57930) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.