

## SLC25A2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57932

## **Product Information**

**Application** IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Human
Host
Clonality
Polyclonal
Calculated MW
32580

## **Additional Information**

**Gene ID** 83884

Other Names Mitochondrial ornithine transporter 2, Solute carrier family 25 member 2,

SLC25A2, ORNT2

**Dilution** IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Protein Information**

Name SLC25A2 ( <u>HGNC:22921</u>)

**Function** Mitochondrial transporter of the positively charged amino acids ornithine,

lysine and arginine, and the neutral amino acid citrulline (PubMed:12807890). In addition, transports the basic amino acids histidine, homoarginine, and asymmetric dimethylarginine (aDMA), but not symmetric DMA, and the D-forms of lysine, arginine, ornithine and histidine (PubMed:12807890, PubMed:26403849). Functions by both counter-exchange and uniport

mechanisms (PubMed:26403849).

**Cellular Location** Mitochondrion membrane; Multi-pass membrane protein. Mitochondrion

inner membrane; Multi-pass membrane protein

**Tissue Location** Expressed in liver, testis, spleen, lung, pancreas, and small intestine and

expressed poorly in other tissues

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.