

SLC25A33 Rabbit pAb

SLC25A33 Rabbit pAb

Catalog # AP57937

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9BSK2
Reactivity	Rat, Mouse
Predicted	Pig, Human, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35375
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SLC25A33
Epitope Specificity	81-180/321
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
SUBCELLULAR LOCATION	Mitochondrion inner membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	SLC25A33 belongs to the SLC25 family of mitochondrial carrier proteins (Haitina et al., 2006 [PubMed 16949250]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	84275
Other Names	Solute carrier family 25 member 33, Bone marrow stromal cell mitochondrial carrier protein, BMSC-MCP, HuBMSC-MCP, Protein PNC1, SLC25A33
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SLC25A33
Function	Mitochondrial transporter that imports/exports pyrimidine nucleotides into and from mitochondria. Selectively transports uridine, thymidine, guanosine, cytosine and inosine (deoxy)nucleoside di- and triphosphates by an antiport

mechanism (PubMed:[25320081](#)). May import (deoxy)nucleoside triphosphates in exchange for intramitochondrial (deoxy)nucleoside diphosphates, thus providing precursors necessary for de novo synthesis of mitochondrial DNA and RNA while exporting products of their catabolism (PubMed:[25320081](#)). Participates in mitochondrial genome maintenance, regulation of mitochondrial membrane potential and mitochondrial respiration (PubMed:[20453889](#)). Upon INS or IGF1 stimulation regulates cell growth and proliferation by controlling mitochondrial DNA replication and transcription, the ratio of mitochondria-to nuclear-encoded components of the electron transport chain resulting in control of mitochondrial ROS production (PubMed:[17596519](#), PubMed:[20453889](#)). Participates in dendritic cell endocytosis and may associate with mitochondrial oxidative phosphorylation (PubMed:[14715278](#)).

Cellular Location

Mitochondrion inner membrane; Multi-pass membrane protein


Tissue Location

Expressed in the central nervous system. Also expressed in testis and skeletal muscle. Weakly expressed in heart, liver, kidney, prostate, colon and peripheral blood leukocytes

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLC25A33) Polyclonal Antibody, Unconjugated (AP57937) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.