

# SLC29A3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57946

## Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">Q9BZD2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51815

## Additional Information

Gene ID	55315
Other Names	Equilibrative nucleoside transporter 3, hENT3, Solute carrier family 29 member 3, SLC29A3, ENT3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

Name	SLC29A3 ( <a href="#">HGNC:23096</a> )
Synonyms	ENT3
Function	Uniporter that mediates the facilitative transport of nucleoside across lysosomal and mitochondrial membranes (PubMed: <a href="#">15701636</a> , PubMed: <a href="#">19164483</a> , PubMed: <a href="#">20595384</a> , PubMed: <a href="#">28729424</a> ). Functions as a non-electrogenic Na(+)-independent transporter (PubMed: <a href="#">15701636</a> , PubMed: <a href="#">19164483</a> , PubMed: <a href="#">28729424</a> ). Substrate transport is pH-dependent and enhanced under acidic condition, probably reflecting the location of the transporter in acidic intracellular compartments (PubMed: <a href="#">15701636</a> , PubMed: <a href="#">19164483</a> , PubMed: <a href="#">28729424</a> ). Proton is not a cotransporting ion but most likely change the ionization state of the transporter which dictates transport- permissible/impermissible conformation for nucleoside translocation (PubMed: <a href="#">28729424</a> ). May direct the nucleoside transport from lysosomes to cytosol or cytosol to mitochondria to facilitate the fundamental function of salvage synthesis of nucleic acids (PubMed: <a href="#">28729424</a> ). Involved in

the transport of nucleosides (adenosine, guanosine, uridine, thymidine, cytidine and inosine) and deoxynucleosides (deoxyadenosine, deoxycytidine) (PubMed:[15701636](#), PubMed:[19164483](#), PubMed:[20595384](#), PubMed:[28729424](#)). Also mediates transport of purine nucleobases (adenine, guanine) and pyrimidine nucleobases (uracil) (PubMed:[15701636](#), PubMed:[19164483](#)). Also able to transport monoamine neurotransmitters dopamine, serotonin, noradrenaline and tyramine (PubMed:[19164483](#)). Capable of transporting ATP (PubMed:[19164483](#)). Mediates nucleoside export from lysosomes in macrophages, which regulates macrophage functions and numbers (By similarity).

**Cellular Location**

Lysosome membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Mitochondrion membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Observed in a punctate intracellular pattern showing partial colocalization with late endosomes/lysosomes (PubMed:15701636). Detected at the cell surface only in certain placental cells (PubMed:19164483)

**Tissue Location**

Widely expressed in both adult and fetal tissues (PubMed:15701636). Highest levels in placenta, uterus, ovary, spleen, lymph node and bone marrow (PubMed:15701636). Expressed in liver (PubMed:19164483). Lowest levels in brain and heart (PubMed:15701636) Expressed in macrophages (PubMed:22174130)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.