

KIF14 Rabbit pAb

KIF14 Rabbit pAb
Catalog # AP58031

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q15058
Reactivity	Rat
Predicted	Dog, Human, Mouse, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	186492
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KIF14
Epitope Specificity	455-550/1648
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle. Note=Nuclear localization observed during interphase in PubMed:16431929 or triggered by entry into mitosis in PubMed:16648480. Cytoplasmic in interphase (PubMed:16648480) and metaphase cells (PubMed:16431929). From prophase to metaphase, accumulates at the developing spindle poles and their associated microtubules. During anaphase, accumulates at the spindle midzone. Localization to the central spindle and midbody during anaphase is dependent upon PRC1 and CIT presence. In cells ready to undergo abscission, concentrates at the contractile ring.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	KIF14 is a member of the kinesin superfamily of microtubule-associated motors (see MIM 148760) that play important roles in intracellular transport and cell division (Nakagawa et al., 1997 [PubMed 9275178]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	9928
Other Names	Kinesin-like protein KIF14, KIF14 (HGNC:19181)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	KIF14 (HGNC:19181)
Function	Microtubule motor protein that binds to microtubules with high affinity through each tubulin heterodimer and has an ATPase activity (By similarity). Plays a role in many processes like cell division, cytokinesis and also in cell proliferation and apoptosis (PubMed:16648480, PubMed:24784001). During cytokinesis, targets to central spindle and midbody through its interaction with PRC1 and CIT respectively (PubMed:16431929). Regulates cell growth through regulation of cell cycle progression and cytokinesis (PubMed:24854087). During cell cycle progression acts through SCF-dependent proteasomal ubiquitin-dependent protein catabolic process which controls CDKN1B degradation, resulting in positive regulation of cyclins, including CCNE1, CCND1 and CCNB1 (PubMed:24854087). During late neurogenesis, regulates the cerebellar, cerebral cortex and olfactory bulb development through regulation of apoptosis, cell proliferation and cell division (By similarity). Also is required for chromosome congression and alignment during mitotic cell cycle process (PubMed:15843429). Regulates cell spreading, focal adhesion dynamics, and cell migration through its interaction with RADIL resulting in regulation of RAP1A-mediated inside-out integrin activation by tethering RADIL on microtubules (PubMed:23209302).
Cellular Location	Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle. Midbody. Note=Nuclear localization observed during interphase (PubMed:16431929). Nuclear localization triggered by entry into mitosis (PubMed:16648480) Cytoplasmic in interphase (PubMed:16648480). Cytoplasmic in metaphase cells (PubMed:16431929). From prophase to metaphase, accumulates at the developing spindle poles and their associated microtubules. During anaphase, accumulates at the spindle midzone. Localization to the central spindle and midbody during anaphase is dependent upon PRC1 and CIT presence. In cells ready to undergo abscission, concentrates at the contractile ring.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF14) Polyclonal Antibody, Unconjugated (AP58031) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.