

Ube2B Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58138

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P63146
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17312
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Ube2B
Epitope Specificity	1-100/152
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane (By similarity). Nucleus (By similarity). Note=In peripheral neurons, expressed both at the plasma membrane and in nuclei (By similarity).
SIMILARITY	Belongs to the ubiquitin-conjugating enzyme family.
SUBUNIT	Interacts with RAD18, UBR2 and WAC.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In association with the E3 enzyme BRE1 (RNF20 and/or RNF40), it plays a role in transcription regulation by catalyzing the monoubiquitination of histone H2B at 'Lys-120' to form H2BK120ub1. H2BK120ub1 gives a specific tag for epigenetic transcriptional activation, elongation by RNA polymerase II, telomeric silencing, and is also a prerequisite for H3K4me and H3K79me formation. In vitro catalyzes 'Lys-11'-, as well as 'Lys-48'- and 'Lys-63'-linked polyubiquitination. Required for postreplication repair of UV-damaged DNA. Associates to the E3 ligase RAD18 to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA-associated PCNA on 'Lys-164'. May be involved in neurite outgrowth.

Additional Information


Gene ID	7320
Other Names	Ubiquitin-conjugating enzyme E2 B, 2.3.2.23, E2 ubiquitin-conjugating enzyme B, RAD6 homolog B, HR6B, hHR6B, Ubiquitin carrier protein B, Ubiquitin-conjugating enzyme E2-17 kDa, Ubiquitin-protein ligase B, UBE2B, RAD6B

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1 µg/Test,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	UBE2B (HGNC:12473)
Function	E2 ubiquitin-conjugating enzyme that accepts ubiquitin from the ubiquitin-activating enzyme E1 and transfers it to a E3 ubiquitin- protein ligase (PubMed: 16337599 , PubMed: 17108083 , PubMed: 17130289 , PubMed: 1717990 , PubMed: 20061386). In vitro catalyzes 'Lys-11'-, as well as 'Lys-48'- and 'Lys-63'-linked polyubiquitination (PubMed: 20061386). Together with the E3 enzyme BRE1 (RNF20 and/or RNF40), plays a role in transcription regulation by catalyzing the monoubiquitination of histone H2B at 'Lys-120' to form H2BK120ub1 (PubMed: 16337599). H2BK120ub1 gives a specific tag for epigenetic transcriptional activation, elongation by RNA polymerase II, telomeric silencing, and is also a prerequisite for H3K4me and H3K79me formation (PubMed: 16337599). May play a role in DNA repair (PubMed: 8062904). Associates to the E3 ligase RAD18 to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA-associated PCNA on 'Lys-164' (PubMed: 17108083 , PubMed: 17130289). In association with the E3 enzyme UBR4, is involved in N-end rule-dependent protein degradation (PubMed: 38182926). May be involved in neurite outgrowth (By similarity).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P63149}. Nucleus {ECO:0000250 UniProtKB:P63149}. Note=In peripheral neurons, expressed both at the plasma membrane and in nuclei {ECO:0000250 UniProtKB:P63149}

Images



Blank control: Hela(blue), the cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice..

Isotype Control Antibody: Rabbit IgG(orange) ;
Secondary Antibody: Goat anti-rabbit IgG-FITC(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA ;
Primary Antibody Dilution: 1 µg in 100 µL1X PBS containing 0.5% BSA(green).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.