

5HT3B Receptor Rabbit pAb

5HT3B Receptor Rabbit pAb Catalog # AP58148

Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession

Reactivity

Host

Clonality

Calculated MW

Physical State

C95264

Human

Rabbit

Polyclonal

50292

Liquid

Immunogen KLH conjugated synthetic peptide derived from human 5HTR3B/5HT3B

receptor

Epitope Specificity 101-200/441

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane; Multi-pass membrane protein. Note=Presumably retained

within the endoplasmic reticulum unless complexed with HTR3A.

SIMILARITY Belongs to the ligand-gated ion channel (TC 1.A.9) family.

5-hydroxytryptamine receptor (TC 1.A.9.2) subfamily. HTR3B sub-subfamily.

SUBUNIT Forms a pentaheteromeric complex with HTR3A. Not functional as a

homomeric complex.

Post-translational modifications Important Note

N-glycosylation required for membrane localization.

This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This is one of the several different receptors for 5-hydroxytryptamine

(serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor is a ligand-gated ion channel, which when activated causes fast, depolarizing responses. It is a cation-specific, but

otherwise relatively nonselective, ion channel.

Additional Information

Gene ID 9177

Other Names 5-hydroxytryptamine receptor 3B, 5-HT3-B, 5-HT3B, Serotonin receptor 3B,

HTR3B (HGNC:5298)

Target/Specificity Expressed in the brain cortex, in the caudate nucleus, the hyppocampus, the

thalamus and the amygdala. Detected in the kidney and testis as well as in monocytes of the spleen, small and large intestine, uterus, prostate, ovary

and placenta.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1ug

/Test

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name HTR3B (HGNC:5298)

Function Forms serotonin (5-hydroxytryptamine/5-HT3)-activated cation- selective

channel complexes, which when activated cause fast, depolarizing responses

in neurons.

Cellular Location Postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane;

Multi-pass membrane protein. Note=Presumably retained within the

endoplasmic reticulum unless complexed with HTR3A

Tissue Location Expressed in the brain cortex, in the caudate nucleus, the hippocampus, the


thalamus and the amygdala. Detected in the kidney and testis as well as in monocytes of the spleen, small and large intestine, uterus, prostate, ovary

and placenta

Background

This is one of the several different receptors for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor is a ligand-gated ion channel, which when activated causes fast, depolarizing responses. It is a cation-specific, but otherwise relatively nonselective, ion channel.

Images




Blank control(blue):293T Cells(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice). Primary Antibody: Rabbit Anti-5HT3B receptor Phospho-c-Fos (Thr325)/AF488 Conjugated antibody (AP58148-AF488), Dilution: 1ug in 100 ul 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG/AF488(orange), used under the same conditions.


Sample: 293T Cell (Human) Lysate at 40 ug

Primary: Anti- 5HT3B receptor (AP58148) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000


dilution

Predicted band size: 50 kD Observed band size: 55 kD





Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (5HT3B receptor) Polyclonal Antibody, Unconjugated (AP58148) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control(blue):293T Cells(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice).
Primary Antibody: Rabbit Anti-5HT3B receptor Phospho-c-Fos (Thr325)/AF488 Conjugated antibody (AP58148-AF488), Dilution: 1ug in 100 ul 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG/AF488(orange) ,used under the same conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.