

ZNF710 Rabbit pAb

ZNF710 Rabbit pAb

Catalog # AP58174

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8N1W2
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74461
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ZNF710
Epitope Specificity	401-500/664
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ZNF710 (Zinc Finger Protein 710) is a Protein Coding gene. Among its related pathways are Gene Expression. GO annotations related to this gene include nucleic acid binding and transcription cofactor activity. An important paralog of this gene is ZNF366.

Additional Information

Gene ID	374655
Other Names	Zinc finger protein 710, ZNF710
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ZNF710
Function	May be involved in transcriptional regulation.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (ZNF710) polyclonal Antibody, Unconjugated (AP58174) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Paraformaldehyde-fixed, paraffin embedded (mouse colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ZNF710) Polyclonal Antibody, Unconjugated (AP58174) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.