

# SEC31A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58214

### **Product Information**

**Application** IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>094979</u>

**Reactivity** Rat, Pig, Dog, Bovine

HostRabbitClonalityPolyclonalCalculated MW133015

## **Additional Information**

**Gene ID** 22872

Other Names Protein transport protein Sec31A, ABP125, ABP130, SEC31-like protein 1,

SEC31-related protein A, Web1-like protein, SEC31A, KIAA0905, SEC31L1

**Dilution** IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

#### **Protein Information**

Name SEC31A

Synonyms KIAA0905, SEC31L1

**Function** Component of the coat protein complex II (COPII) which promotes the

formation of transport vesicles from the endoplasmic reticulum (ER) (PubMed:10788476). The coat has two main functions, the physical

deformation of the endoplasmic reticulum membrane into vesicles and the

selection of cargo molecules (By similarity).

**Cellular Location** Cytoplasmic vesicle, COPII-coated vesicle membrane; Peripheral

membrane protein; Cytoplasmic side. Endoplasmic reticulum membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=Associates with membranes in a GTP- dependent manner (By similarity). Localizes to endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:17428803, PubMed:25201882,

PubMed:28442536) {ECO:0000250 | UniProtKB:Q9Z2Q1,

ECO:0000269 | PubMed:17428803, ECO:0000269 | PubMed:25201882, ECO:0000269 | PubMed:28442536}

## **Tissue Location**

Abundantly and ubiquitously expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.