

PMVK Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58293

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q15126
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21995
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PMVK
Epitope Specificity	15-120/192
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Peroxisome.
SUBUNIT	Monomer (By similarity).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	PMVK is a 192 amino acid peroxisomal enzyme belonging to the nucleoside monophosphate (NMP) kinase family and is expressed in heart, liver, skeletal muscle, kidney, and pancreas with lower expression in brain, placenta and lung. Induced by sterol, PMVK participates in isopentenyl diphosphate biosynthesis via the mevalonate pathway. PMVK catalyzes the conversion of mevalonate 5-phosphate into mevalonate 5-diphosphate in the fifth reaction of the cholesterol biosynthetic pathway. PMVK exists as a monomer and is encoded by a gene located on human chromosome 1, which houses over 3,000 genes and is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome.

Additional Information

Gene ID	10654
Other Names	Phosphomevalonate kinase, PMKase, hPMK, 2.7.4.2, PMVK, PMKI
Target/Specificity	Heart, liver, skeletal muscle, kidney, and pancreas. Lower level in brain, placenta and lung.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce


Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	PMVK
Synonyms	PMKI
Function	Catalyzes the reversible ATP-dependent phosphorylation of mevalonate 5-phosphate to produce mevalonate diphosphate and ADP, a key step in the mevalonic acid mediated biosynthesis of isopentenyl diphosphate and other polyisoprenoid metabolites.
Cellular Location	Cytoplasm, cytosol
Tissue Location	Heart, liver, skeletal muscle, kidney, and pancreas. Lower level in brain, placenta and lung

Images



Tissue/cell: human kidney tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal
goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-PMVK Polyclonal Antibody,
Unconjugated(AP58293) 1:200, overnight at 4°C, followed
by conjugation to the secondary antibody(SP-0023) and
DAB(C-0010) staining



Sample:
Uterus (Mouse) Lysate at 40 ug
Cerebrum (Mouse) Lysate at 40 ug
Heart (Mouse) Lysate at 40 ug
Lymph node (Mouse) Lysate at 40 ug
293T(Human) Cell Lysate at 30 ug
Primary: Anti-PMVK (AP58293) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution
Predicted band size: 22 kD
Observed band size: 24 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.