

RAIDD Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58318

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P78560
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22745
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RAIDD
Epitope Specificity	31-130/199
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 CARD domain.Contains 1 death domain.
SUBUNIT	Interacts with LRDD.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Apoptotic adaptor molecule specific for caspase-2 and FASL/TNF receptor-interacting protein RIP. In the presence of RIP and TRADD, CRADD recruits caspase-2 to the TNFR-1 signalling complex.


Additional Information

Gene ID	8738
Other Names	Death domain-containing protein CRADD, Caspase and RIP adapter with death domain, RIP-associated protein with a death domain, CRADD, RAIDD
Target/Specificity	Constitutively expressed in most tissues, with particularly high expression in adult heart, testis, liver, skeletal muscle, fetal liver and kidney.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	CRADD
Synonyms	RAIDD
Function	Adapter protein that associates with PIDD1 and the caspase CASP2 to form the PIDDosome, a complex that activates CASP2 and triggers apoptosis (PubMed: 15073321 , PubMed: 16652156 , PubMed: 17159900 , PubMed: 17289572 , PubMed: 9044836). Also recruits CASP2 to the TNFR-1 signaling complex through its interaction with RIPK1 and TRADD and may play a role in the tumor necrosis factor-mediated signaling pathway (PubMed: 8985253).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:O88843}. Nucleus {ECO:0000250 UniProtKB:O88843}
Tissue Location	Constitutively expressed in most tissues, with particularly high expression in adult heart, testis, liver, skeletal muscle, fetal liver and kidney.

Images



Sample:
Lymph node (Mouse) Lysate at 40 ug
Primary: Anti-RAIDD (AP58318) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 22 kD
Observed band size: 22 kD



Sample:
Lymph node (Mouse) Lysate at 40 ug
Primary: Anti- RAIDD (AP58318) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 22 kD
Observed band size: 22 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.