

# SMAGP Rabbit pAb

SMAGP Rabbit pAb

Catalog # AP58331

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, IF, E
<b>Primary Accession</b>	<a href="#">Q0VAQ4</a>
<b>Predicted</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	10679
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human SMAGP
<b>Epitope Specificity</b>	24-80/97
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Cell membrane; Single-pass type III membrane protein. Cytoplasmic vesicle membrane; Single-pass type III membrane protein. Note=Predominantly on lateral parts of the membrane, at cell-cell epithelial junctions. Detected on cytoplasmic membranes in undifferentiated tumors.
<b>SIMILARITY</b>	Belongs to the SMAGP family.
<b>Post-translational modifications</b>	O-glycosylated. The O-glycan is modified with sialic acid residues.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## Additional Information

---

<b>Gene ID</b>	57228
<b>Other Names</b>	Small cell adhesion glycoprotein, Small transmembrane and glycosylated protein, SMAGP
<b>Target/Specificity</b>	Detected in breast, endometrium, colon and biliary tract. Detected in polarized epithelial structures characterized by cell-cell adhesion (at protein level).
<b>Dilution</b>	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	SMAGP
<b>Function</b>	May play a role in epithelial cell-cell contacts. May play a role in tumor invasiveness and metastasis formation.
<b>Cellular Location</b>	Cell membrane; Single-pass type III membrane protein. Cytoplasmic vesicle membrane; Single-pass type III membrane protein. Note=Predominantly on lateral parts of the membrane, at cell-cell epithelial junctions (PubMed:15021913) Detected on cytoplasmic membranes in undifferentiated tumors (PubMed:15986429).
<b>Tissue Location</b>	Detected in breast, endometrium, colon and biliary tract. Detected in polarized epithelial structures characterized by cell-cell adhesion (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.