

ADAM18 Rabbit pAb

ADAM18 Rabbit pAb
Catalog # AP58332

Product Information

Application	WB
Reactivity	Mouse
Predicted	Human, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ADAM18
Epitope Specificity	301-400/739
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
SIMILARITY	Contains 1 disintegrin domain. Contains 1 EGF-like domain. Contains 1 peptidase M12B domain.
Post-translational modifications	The prodomain and the metalloprotease-like domain are cleaved during the epididymal maturation of the spermatozoa.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ADAM18 was first described as tMDC III (testis-specific Metalloproteinase-like disintegrin-like cysteine-rich protein-III). ADAM18 is a member of the ADAMs family (A Disintegrin And Metalloproteinase), but does not contain the canonical HExxHxxxxH zinc-binding metalloproteinase catalytic site. The domain structure of the full-length ADAM18 includes a signal sequence, propeptide domain, metalloproteinase-like domain, disintegrin-like domain, cys-rich domain, EGF-like domain, a short spacer region, then the transmembrane domain and a cytoplasmic domain. Like many of the reproductive-specific ADAMs, ADAM18 plays a nonenzymatic role, or (as is the case for ADAMs 1 & 2 (fertilin alpha and beta)), the protein acts in concert with a proteolytically active ADAM to process proteins. Little is known about interactions between ADAM18 and other ADAMs. To date, 4 different sequences for human ADAM18 are published; 739, 715, 182 and 153 amino acids in length.

Additional Information

Other Names	Disintegrin and metalloproteinase domain-containing protein 18, ADAM 18, Transmembrane metalloproteinase-like, disintegrin-like, and cysteine-rich protein III, tMDC III, ADAM18, TMDC3
Target/Specificity	Expressed specifically in testis.


Dilution WB=1:500-2000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

ADAM18 was first described as tMDC III (testis-specific Metalloproteinase-like disintegrin-like cysteine-rich protein-III). ADAM18 is a member of the ADAMs family (A Disintegrin And Metalloproteinase), but does not contain the canonical HExxHxxxxH zinc-binding metalloproteinase catalytic site. The domain structure of the full-length ADAM18 includes a signal sequence, propeptide domain, metalloproteinase-like domain, disintegrin-like domain, cys-rich domain, EGF-like domain, a short spacer region, then the transmembrane domain and a cytoplasmic domain. Like many of the reproductive-specific ADAMs, ADAM18 plays a nonenzymatic role, or (as is the case for ADAMs 1 & 2 (fertilin alpha and beta)), the protein acts in concert with a proteolytically active ADAM to process proteins. Little is known about interactions between ADAM18 and other ADAMs. To date, 4 different sequences for human ADAM18 are published; 739, 715, 182 and 153 amino acids in length.

Images



Sample:

Heart (Mouse) Lysate at 40 ug

Primary: Anti- ADAM18 (AP58332) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 62 kD

Observed band size: 75/50 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.