

MAS1L Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58347

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat
Host
Clonality
Polyclonal
Calculated MW
Physical State
P35410
Rat
Polyclonal
42411
Liquid

Immunogen KLH conjugated synthetic peptide derived from human MAS1L

Epitope Specificity 201-300/378

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane; Multi-pass membrane protein.

SIMILARITY Belongs to the G-protein coupled receptor 1 family. Mas subfamily. **Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions Belongs to the G-protein coupled receptor 1 family. Mas subfamily.

Additional Information

Gene ID 116511

Other Names Mas-related G-protein coupled receptor MRG, MAS-R, MAS1-like, MAS1L, MRG

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information


Name MAS1L

Synonyms MRG

Cellular Location Cell membrane; Multi-pass membrane protein.


Images






Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-MAS1L Polyclonal Antibody, Unconjugated(AP58347) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human lung carcinoma; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min;

Incubation: Anti-MAS1L Polyclonal Antibody, Unconjugated(AP58347) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MAS1L) Polyclonal Antibody, Unconjugated (AP58347) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.