

ABI2 Rabbit pAb

ABI2 Rabbit pAb
Catalog # AP58354

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9NYB9
Reactivity	Mouse, Rat
Predicted	Human, Chicken, Dog, Pig, Horse, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55663
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ABI2
Epitope Specificity	421-513/513
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm (By similarity). Isoform 1: Cell projection, lamellipodium (By similarity). Cell projection, filopodium (By similarity). Cytoplasm, cytoskeleton (By similarity). Note=Isoform 1 but not isoform 3 is localized to protruding lamellipodia and filopodia tips (By similarity). Belongs to the ABI family. Contains 1 SH3 domain. Contains 1 t-SNARE coiled-coil homology domain. Is a substrate for ABL1.
SIMILARITY	
Post-translational modifications	
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ABI2 (Abelson interactor 2) belongs to the ABI family. The Abelson interactors are among a group of proteins that interact with Abl tyrosine kinase, a non-receptor tyrosine kinase whose activation results in the regulation of cytoskeleton function, cell growth and apoptosis in response to a variety of biological stimuli.

Additional Information

Gene ID	10152
Other Names	Abl interactor 2 {ECO:0000303 Ref.16}, Abelson interactor 2 {ECO:0000303 Ref.16}, Abi-2, Abl-binding protein 3, AblBP3, Arg-binding protein 1, ArgBP1, ABI2 {ECO:0000303 PubMed:28397838, ECO:0000312 HGNC:HGNC:24011}
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ABI2 {ECO:0000303 PubMed:28397838, ECO:0000312 HGNC:HGNC:24011}
Function	Regulator of actin cytoskeleton dynamics underlying cell motility and adhesion. Functions as a component of the WAVE complex, which activates actin nucleating machinery Arp2/3 to drive lamellipodia formation (PubMed: 21107423). Acts as a regulator and substrate of nonreceptor tyrosine kinases ABL1 and ABL2 involved in processes linked to cell growth and differentiation. Positively regulates ABL1-mediated phosphorylation of ENAH, which is required for proper polymerization of nucleated actin filaments at the leading edge (PubMed: 10498863 , PubMed: 7590236 , PubMed: 8649853). Contributes to the regulation of actin assembly at the tips of neuron projections. In particular, controls dendritic spine morphogenesis and may promote dendritic spine specification toward large mushroom-type spines known as repositories of memory in the brain (By similarity). In hippocampal neurons, may mediate actin-dependent BDNF-NTRK2 early endocytic trafficking that triggers dendrite outgrowth (By similarity). Participates in ocular lens morphogenesis, likely by regulating lamellipodia-driven adherens junction formation at the epithelial cell-secondary lens fiber interface (By similarity). Also required for nascent adherens junction assembly in epithelial cells (PubMed: 15572692).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed. Abundant in testes, ovary, thymus, and colon, with lower but detectable levels in prostate, peripheral blood leukocytes, and spleen.

Background

ABI2 (Abelson interactor 2) belongs to the ABI family. The Abelson interactors are among a group of proteins that interact with Abl tyrosine kinase, a non-receptor tyrosine kinase whose activation results in the regulation of cytoskeleton function, cell growth and apoptosis in response to a variety of biological stimuli.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ABI2) Polyclonal Antibody, Unconjugated (AP58354) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.