

CD66c Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58364

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P40199
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37237
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CD66c/CEACAM6
Epitope Specificity	165-260/344
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Lipid-anchor, GPI-anchor.
SIMILARITY	Belongs to the immunoglobulin superfamily. CEA family. Contains 2 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
SUBUNIT	Homodimer. Binding of E.coli Dr adhesins leads to dissociation of the homodimer.
Post-translational modifications	Complex immunoreactive glycoprotein with a MW of 180 kDa comprising 60% carbohydrate.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	CEA-related cell adhesion molecules (CEACAM) belong to the carcinoembryonic antigen (CEA) family. It consists of seven CEACAM (CEACAM 1, CEACAM 3-CEACAM 8) and 11 pregnancy-specific glyco-protein (PSG 1-PSG 11) members. The CEA family proteins belong to the immunoglobulin (Ig) superfamily and are composed of one Ig variable-like (IgV) and a varying number (0-6) of Ig constant-like (IgC) domains. CEACAM molecules are membrane-bound either via a transmembrane domain or a glycosyl phosphatidyl inositol (GPI) anchor. CEACAM molecules are differentially expressed in epithelial cells or in leucocytes. Over-expression of CEA/CEACAM 5 in tumors of epithelial origin is the basis of its wide-spread use as a tumor marker. The function of CEACAM family members varies widely: they function as cell adhesion molecules, tumor suppressors, regulators of lymphocyte and dendritic cell activation, receptors of Neisseria species and other bacteria.

Additional Information

Gene ID 4680

Other Names	Carcinoembryonic antigen-related cell adhesion molecule 6, Non-specific crossreacting antigen, Normal cross-reacting antigen, CD66c, CEACAM6, NCA
Target/Specificity	Found in adenocarcinomas of endodermally derived digestive system epithelium and fetal colon.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CEACAM6 (HGNC:1818)
Function	Cell surface glycoprotein that plays a role in cell adhesion and tumor progression (PubMed: 10910050 , PubMed: 11590190 , PubMed: 1378450 , PubMed: 16204051 , PubMed: 2022629 , PubMed: 2803308 , PubMed: 8776764). Intercellular adhesion occurs in a calcium- and fibronectin-independent manner (PubMed: 16204051 , PubMed: 2022629). Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM5 and CEACAM8 (PubMed: 11590190 , PubMed: 16204051 , PubMed: 2022629 , PubMed: 2803308 , PubMed: 8776764). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed: 8776764). Plays a role in neutrophil adhesion to cytokine-activated endothelial cells (PubMed: 1378450). Plays a role in cell migration and cell adhesion to endothelial cells (PubMed: 16204051).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Apical cell membrane. Cell surface. Note=Localized to the apical glycocalyx surface.
Tissue Location	Expressed in neutrophils (PubMed:1378450). Expressed in columnar epithelial and goblet cells of the colon (PubMed:10436421). Expressed in numerous tumor cell lines (at protein level) (PubMed:16204051).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.