

EVC2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58482

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q86UK5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	147948

Additional Information

Gene ID	132884
Other Names	Limbin, Ellis-van Creveld syndrome protein 2, EVC2, EVC2, LBN
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EVC2
Synonyms	LBN
Function	Component of the EvC complex that positively regulates ciliary Hedgehog (Hh) signaling. Plays a critical role in bone formation and skeletal development. May be involved in early embryonic morphogenesis.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:Q8K1G2}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:Q8K1G2} Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250 UniProtKB:Q8K1G2}. Cell projection, cilium {ECO:0000250 UniProtKB:Q8K1G2}. Cell projection, cilium membrane {ECO:0000250 UniProtKB:Q8K1G2}. Nucleus {ECO:0000250 UniProtKB:Q8K1G2} Note=The EvC complex localizes at the base of cilia in the EvC zone of primary cilia in a EFCAB7-dependent manner {ECO:0000250 UniProtKB:Q8K1G2}
Tissue Location	Found in the heart, placenta, lung, liver, skeletal muscle, kidney and pancreas.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.