

CRMP1 Rabbit pAb

CRMP1 Rabbit pAb
Catalog # AP58483

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q14194
Predicted	Human, Mouse, Rat, Chicken, Dog, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62184
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CRMP1
Epitope Specificity	121-220/572
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Cytoplasm, cytoskeleton, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Associated with centrosomes and the mitotic spindle during metaphase.
SIMILARITY	Belongs to the DHOase family. Hydantoinase/dihydropyrimidinase subfamily.
SUBUNIT	Homotetramer, and heterotetramer with DPYSL2, DPYSL3, DPYSL4 or DPYSL5. Interacts with PLXNA1 (By similarity).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of a family of cytosolic phosphoproteins expressed exclusively in the nervous system. The encoded protein is thought to be a part of the semaphorin signal transduction pathway implicated in semaphorin-induced growth cone collapse during neural development. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	1400
Other Names	Dihydropyrimidinase-related protein 1, DRP-1, Collapsin response mediator protein 1, CRMP-1, Inactive dihydropyrimidinase, Unc-33-like phosphoprotein 3, ULIP-3, CRMP1, DPYSL1, ULIP3
Target/Specificity	Brain.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CRMP1
Synonyms	DPYSL1, ULIP3
Function	Necessary for signaling by class 3 semaphorins and subsequent remodeling of the cytoskeleton (PubMed: 25358863). Plays a role in axon guidance (PubMed: 25358863). During the axon guidance process, acts downstream of SEMA3A to promote FLNA dissociation from F-actin which results in the rearrangement of the actin cytoskeleton and the collapse of the growth cone (PubMed: 25358863). Involved in invasive growth and cell migration (PubMed: 11562390). May participate in cytokinesis (PubMed: 19799413).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Cell projection, growth cone {ECO:0000250 UniProtKB:P97427}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P97427}. Perikaryon {ECO:0000250 UniProtKB:P97427}. Note=Associated with centrosomes and the mitotic spindle during metaphase (PubMed:11562390). Colocalizes with FLNA and tubulin in the central region of DRG neuron growth cone (By similarity). Following SEMA3A stimulation of DRG neurons, colocalizes with F-actin (By similarity) {ECO:0000250 UniProtKB:P97427, ECO:0000269 PubMed:11562390}
Tissue Location	Brain.

Background

This gene encodes a member of a family of cytosolic phosphoproteins expressed exclusively in the nervous system. The encoded protein is thought to be a part of the semaphorin signal transduction pathway implicated in semaphorin-induced growth cone collapse during neural development. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2008].

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.