

# NLRP10 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58532

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF, E
<b>Primary Accession</b>	<a href="#">Q86W26</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	75032

## Additional Information

---

<b>Gene ID</b>	338322
<b>Other Names</b>	NACHT, LRR and PYD domains-containing protein 10, Nucleotide-binding oligomerization domain protein 8, NLRP10, NALP10, NOD8, PYNOD
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	NLRP10
<b>Synonyms</b>	NALP10, NOD8, PYNOD
<b>Function</b>	Inhibits autoprocessing of CASP1, CASP1-dependent IL1B secretion, PYCARD aggregation and PYCARD-mediated apoptosis but not apoptosis induced by FAS or BID (PubMed: <a href="#">15096476</a> ). Displays anti- inflammatory activity (PubMed: <a href="#">20393137</a> ). Required for immunity against C.albicans infection (By similarity). Involved in the innate immune response by contributing to pro-inflammatory cytokine release in response to invasive bacterial infection (PubMed: <a href="#">22672233</a> ). Contributes to T-cell-mediated inflammatory responses in the skin (By similarity). Plays a role in protection against periodontitis through its involvement in induction of IL1A via ERK activation in oral epithelial cells infected with periodontal pathogens (PubMed: <a href="#">28766990</a> ). Exhibits both ATPase and GTPase activities (PubMed: <a href="#">23861819</a> ).
<b>Cellular Location</b>	Cytoplasm. Cell membrane; Peripheral membrane protein. Note=Cytoplasmic protein which is recruited to the cell membrane by NOD1 following invasive bacterial infection

**Tissue Location**

Highly expressed in basal and suprabasal epidermal cell layers with lower levels in dermal fibroblast cells (at protein level) (PubMed:22672233). Widely expressed with highest levels in heart, brain and skeletal muscle (PubMed:15096476). Also expressed in liver, colon, dermis and epidermis (PubMed:15096476). Little expression detected in myeloid cells or peripheral blood mononuclear cells (PubMed:15096476).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.