

NLRP10 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58532

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	<u>Q86W26</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	75032

Additional Information

Gene ID	338322
Other Names	NACHT, LRR and PYD domains-containing protein 10, Nucleotide-binding oligomerization domain protein 8, NLRP10, NALP10, NOD8, PYNOD
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NLRP10
Synonyms	NALP10, NOD8, PYNOD
Function	Inhibits autoprocessing of CASP1, CASP1-dependent IL1B secretion, PYCARD aggregation and PYCARD-mediated apoptosis but not apoptosis induced by FAS or BID (PubMed: <u>15096476</u>). Displays anti- inflammatory activity (PubMed: <u>20393137</u>). Required for immunity against C.albicans infection (By similarity). Involved in the innate immune response by contributing to pro-inflammatory cytokine release in response to invasive bacterial infection (PubMed: <u>22672233</u>). Contributes to T-cell-mediated inflammatory responses in the skin (By similarity). Plays a role in protection against periodontitis through its involvement in induction of IL1A via ERK activation in oral epithelial cells infected with periodontal pathogens (PubMed: <u>28766990</u>). Exhibits both ATPase and GTPase activities (PubMed: <u>23861819</u>).
Cellular Location	Cytoplasm. Cell membrane; Peripheral membrane protein. Note=Cytoplasmic protein which is recruited to the cell membrane by NOD1 following invasive

bacterial infection

Tissue LocationHighly expressed in basal and suprabasal epidermal cell layers with lower
levels in dermal fibroblast cells (at protein level) (PubMed:22672233). Widely
expressed with highest levels in heart, brain and skeletal muscle
(PubMed:15096476). Also expressed in liver, colon, dermis and epidermis
(PubMed:15096476). Little expression detected in myeloid cells or peripheral
blood mononuclear cells (PubMed:15096476).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.