

# SOX10 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5854C

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P56693</a>
<b>Other Accession</b>	<a href="#">Q55170</a> , <a href="#">A5A763</a> , <a href="#">Q04888</a> , <a href="#">NP_008872.1</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Predicted</b>	Rat, Pig
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	49911
<b>Antigen Region</b>	231-259

## Additional Information

---

<b>Gene ID</b>	6663
<b>Other Names</b>	Transcription factor SOX-10, SOX10
<b>Target/Specificity</b>	This SOX10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 231-259 amino acids of human SOX10.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SOX10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	SOX10
<b>Function</b>	Transcription factor that plays a central role in developing and mature glia (By similarity). Specifically activates expression of myelin genes, during oligodendrocyte (OL) maturation, such as DUSP15 and MYRF, thereby playing a central role in oligodendrocyte maturation and CNS myelination (By similarity). Once induced, MYRF cooperates with SOX10 to implement the

myelination program (By similarity). Transcriptional activator of MITF, acting synergistically with PAX3 (PubMed:[21965087](#)). Transcriptional activator of MBP, via binding to the gene promoter (By similarity).

#### Cellular Location


Cytoplasm. Nucleus. Mitochondrion outer membrane {ECO:0000250|UniProtKB:Q04888}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q04888}; Cytoplasmic side {ECO:0000250|UniProtKB:Q04888}

#### Tissue Location


Expressed in fetal brain and in adult brain, heart, small intestine and colon

## Images

---



AP5854c staining SOX10 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-SOX10 Antibody (Center) at 1:2000 dilution  
Lane 1: C6 whole cell lysates Lane 2: A375 whole cell lysates  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 50 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

## Citations

---

- [Modulation of Sox10, HIF-1α, Survivin, and YAP by Minocycline in the Treatment of Neurodevelopmental Handicaps following Hypoxic Insult.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.