10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CCDC50 Rabbit pAb

CCDC50 Rabbit pAb
Catalog # AP58550

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8IVM0
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35822
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CCDC50
Epitope Specificity	251-306/306
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
DISEASE	Defects in CCDC50 are the cause of deafness autosomal dominant type 44 (DFNA44) . A form of non-syndromic hearing loss. It is initially moderate and affects mainly low to mid frequencies. Later, it progresses to involve all the frequencies and leads to a profound hearing loss by the 6th decade. The onset of the hearing loss occurs in the first decade of life.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with multiple ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual, progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the inner ear during development and postnatal maturation and associates with microtubule-based structures. This protein may also function as a negative regulator of NF-kB signaling and as an effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008].

Additional Information

Gene ID	152137
Other Names	Coiled-coil domain-containing protein 50, Protein Ymer, CCDC50, C3orf6
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	CCDC50
Synonyms	C3orf6
Function	Involved in EGFR signaling.
Cellular Location	Cytoplasm. Note=Associated with microtubules of the cytoskeleton and mitotic apparatus.
Tissue Location	Isoform 1 and isoform 2 are coexpressed in placenta, liver, lung, kidney and pancreas. Only isoform 1 is detected in skeletal muscle, brain and heart.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Cerebellum (Mouse) Lysate at 40 ug
Primary: Anti- CCDC50 (AP58550) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 34 kD
Observed band size: 33 kD



Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCDC50) Polyclonal Antibody, Unconjugated (AP58550) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.