

PLSCR3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58592

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9NRY6
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31648

Additional Information

Gene ID	57048
Other Names	Phospholipid scramblase 3, PL scramblase 3, Ca(2+)-dependent phospholipid scramblase 3, PLSCR3
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	PLSCR3
Function	Catalyzes calcium-induced ATP-independent rapid bidirectional and non-specific movement of the phospholipids (lipid scrambling or lipid flip-flop) between the inner and outer membrane of the mitochondria (PubMed: 14573790 , PubMed: 17226776 , PubMed: 18358005 , PubMed: 29337693 , PubMed: 31769662). Plays an important role in mitochondrial respiratory function, morphology, and apoptotic response (PubMed: 12649167 , PubMed: 14573790 , PubMed: 17226776 , PubMed: 18358005). Mediates the translocation of cardiolipin from the mitochondrial inner membrane to outer membrane enhancing t-Bid induced cytochrome c release and apoptosis (PubMed: 14573790 , PubMed: 17226776 , PubMed: 18358005). Enhances TNFSF10-induced apoptosis by regulating the distribution of cardiolipin in the mitochondrial membrane resulting in increased release of apoptogenic factors and consequent amplification of the activity of caspases (PubMed: 18491232). Regulates cardiolipin de novo biosynthesis and its resynthesis (PubMed: 16939411).

Cellular Location	<p>Mitochondrion membrane; Single-pass type II membrane protein {ECO:0000250 UniProtKB:Q6QBB4}. Mitochondrion inner membrane {ECO:0000250 UniProtKB:Q6QBB4}; Single-pass type II membrane protein {ECO:0000250 UniProtKB:Q6QBB4}. Nucleus {ECO:0000250 UniProtKB:Q9JIZ9} Note=Palmitoylation regulates its localization to the cell membrane or the nucleus; trafficking to the cell membrane is dependent upon palmitoylation whereas in the absence of palmitoylation, localizes to the nucleus. {ECO:0000250 UniProtKB:Q9JIZ9}</p>
Tissue Location	<p>Expressed in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, uterus, small intestine and peripheral blood lymphocytes. Not detected in testis, brain and liver</p>

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.