

VKORC1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5859A

Product Information

Application	WB, IHC-P, E
Primary Accession	Q9BQB6
Other Accession	Q6B4J2 , NP_076869.1
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	18235
Antigen Region	15-41

Additional Information

Gene ID	79001
Other Names	Vitamin K epoxide reductase complex subunit 1, Vitamin K1 2, 3-epoxide reductase subunit 1, VKORC1, VKOR
Target/Specificity	This VKORC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-41 amino acids from the N-terminal region of human VKORC1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	VKORC1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	VKORC1 {ECO:0000303 PubMed:14765194, ECO:0000312 HGNC:HGNC:23663}
Function	Involved in vitamin K metabolism. Catalytic subunit of the vitamin K epoxide reductase (VKOR) complex which reduces inactive vitamin K 2,3-epoxide to

active vitamin K. Vitamin K is required for the gamma-carboxylation of various proteins, including clotting factors, and is required for normal blood coagulation, but also for normal bone development.


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein


Tissue Location

Expressed at highest levels in fetal and adult liver, followed by fetal heart, kidney, and lung, adult heart, and pancreas.

Images



Anti-VKORC1 Antibody (N-term) at 1:1000 dilution + Human lung whole tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



VKORC1 Antibody (N-term) (Cat. #AP5859a) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the VKORC1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.