

ARHGEF18 Rabbit pAb

ARHGEF18 Rabbit pAb

Catalog # AP58608

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q6ZSZ5
Predicted	Rat, Pig, Dog, Human, Mouse, Chicken, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	151642
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ARHGEF18/p114RhoGEF
Epitope Specificity	501-600/1173
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Note: Colocalizes with actin stress fibers.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Rho GTPases are GTP binding proteins that regulate a wide spectrum of cellular functions. These cellular processes include cytoskeletal rearrangements, gene transcription, cell growth and motility. Activation of Rho GTPases is under the direct control of guanine nucleotide exchange factors (GEFs). The protein encoded by this gene is a guanine nucleotide exchange factor and belongs to the Rho GTPase GFE family. Family members share a common feature, a Dbl (DH) homology domain followed by a pleckstrin (PH) homology domain. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2008].

Additional Information

Gene ID	23370
Other Names	Rho guanine nucleotide exchange factor 18, 114 kDa Rho-specific guanine nucleotide exchange factor, p114-Rho-GEF, p114RhoGEF, Septin-associated RhoGEF, SA-RhoGEF, ARHGEF18, KIAA0521
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ARHGEF18
Synonyms	KIAA0521
Function	Acts as a guanine nucleotide exchange factor (GEF) for RhoA GTPases. Its activation induces formation of actin stress fibers. Also acts as a GEF for RAC1, inducing production of reactive oxygen species (ROS). Does not act as a GEF for CDC42. The G protein beta-gamma (Gbetagamma) subunits of heterotrimeric G proteins act as activators, explaining the integrated effects of LPA and other G protein-coupled receptor agonists on actin stress fiber formation, cell shape change and ROS production. Required for EPB41L4B-mediated regulation of the circumferential actomyosin belt in epithelial cells (PubMed: 22006950).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton. Cell membrane. Apical cell membrane. Note=In unactivated eosinophils, distributed around the cell periphery in the perimembranous region (PubMed:29601110). In activated eosinophils, relocates to the tip of the nucleopod, a membrane structure formed during activation when the nucleus moves to one end of the cell, and is also concentrated in membrane protrusions at the opposite end of the cell (PubMed:29601110) Localizes to the apical cell membrane in epithelial cells (PubMed:22006950).
Tissue Location	Expressed in all tissues tested with highest expression in kidney and pancreas. Weakly or not expressed in liver, skeletal muscle and testis. Isoform 1: Expressed in eosinophils (PubMed:29601110). Isoform 2: Expressed in eosinophils (PubMed:29601110). Isoform 3: Expressed in eosinophils (PubMed:29601110). Isoform 4: Not detected in eosinophils (PubMed:29601110).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.