

# VAMP4 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5861c

## Product Information

---

|                          |  |
|--------------------------|--|
| <b>Application</b>       | WB, IHC-P, E   |
| <b>Primary Accession</b> | <a href="#">O75379</a>   |
| <b>Other Accession</b>   | <a href="#">O70480</a> , <a href="#">Q32L97</a> , <a href="#">NP_001172056.1</a> , <a href="#">NP_003753.2</a> |
| <b>Reactivity</b>        | Human, Mouse   |
| <b>Predicted</b>         | Bovine   |
| <b>Host</b>              | Rabbit   |
| <b>Clonality</b>         | Polyclonal   |
| <b>Isotype</b>           | Rabbit IgG   |
| <b>Clone Names</b>       | RB22111  |
| <b>Calculated MW</b>     | 16397  |
| <b>Antigen Region</b>    | 16-45  |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 8674   |
| <b>Other Names</b>        | Vesicle-associated membrane protein 4, VAMP-4, VAMP4   |
| <b>Target/Specificity</b> | This VAMP4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 16-45 amino acids from the Central region of human VAMP4.            |
| <b>Dilution</b>           | WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.  |
| <b>Format</b>             | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                      |
| <b>Precautions</b>        | VAMP4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.  |

## Protein Information

---

|                 |   |
|-----------------|---|
| <b>Name</b>     | VAMP4   |
| <b>Function</b> | Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical |


for remodeling the secretory response of granule.

## Cellular Location


Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrane protein Note=Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles

## Images


---



VAMP4 Antibody (Center) (Cat. #AP5861c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the VAMP4 antibody detected the VAMP4 protein (arrow).



VAMP4 Antibody (Center) (Cat. #AP5861c) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the VAMP4 antibody detected the VAMP4 protein (arrow).



VAMP4 antibody (Center) (Cat. #AP5861c) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the VAMP4 antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.