

IQGAP2 Rabbit pAb

IQGAP2 Rabbit pAb Catalog # AP58623

Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession <u>Q13576</u>

Predicted Human, Mouse, Rat, Dog, Horse, Rabbit, Sheep

Host Rabbit
Clonality Polyclonal
Calculated MW 180578
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human IQGAP2

Epitope Specificity 51-150/1575

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SIMILARITY Contains 1 CH (calponin-homology) domain. Contains 3 IQ domains. Contains

1 Ras-GAP domain. Contains 1 WW domain.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This gene encodes a member of the IQGAP family. The protein contains three

IQ domains, one calponin homology domain, one Ras-GAP domain and one WW domain. It interacts with components of the cytoskeleton, with cell adhesion molecules, and with several signaling molecules to regulate cell

morphology and motility. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID 10788

Other Names Ras GTPase-activating-like protein IQGAP2, IQGAP2

Target/Specificity Isoform 2 expression is enhanced in testis.

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500

0-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name IQGAP2

Function Binds to activated CDC42 and RAC1 but does not seem to stimulate their

GTPase activity. Associates with calmodulin.

Tissue Location Isoform 2 expression is enhanced in testis.

Background

This gene encodes a member of the IQGAP family. The protein contains three IQ domains, one calponin homology domain, one Ras-GAP domain and one WW domain. It interacts with components of the cytoskeleton, with cell adhesion molecules, and with several signaling molecules to regulate cell morphology and motility. [provided by RefSeq, Jul 2008]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.