

Tankyrase 2 Rabbit pAb

Tankyrase 2 Rabbit pAb

Catalog # AP58658

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9H2K2
Reactivity	Rat
Predicted	Pig, Human, Mouse, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	126918
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Tankyrase 2
Epitope Specificity	1-100/1166
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Golgi apparatus membrane. Nucleus. Chromosome > telomere. Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles. Also found around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS2 is found in the nucleus, associated with TRF1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	80351
Other Names	Poly [ADP-ribose] polymerase tankyrase-2, 2.4.2.30, ADP-ribosyltransferase diphtheria toxin-like 6, ARTD6, Poly [ADP-ribose] polymerase 5B, Protein poly-ADP-ribosyltransferase tankyrase-2, 2.4.2.-, TNKS-2, TRF1-interacting ankyrin-related ADP-ribose polymerase 2, Tankyrase II, Tankyrase-2, TANK2, Tankyrase-like protein, Tankyrase-related protein, TNKS2 (HGNC:15677)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	TNKS2 (HGNC:15677)
-------------	--------------------------------------

Function	Poly-ADP-ribosyltransferase involved in various processes such as Wnt signaling pathway, telomere length and vesicle trafficking (PubMed:11739745, PubMed:11802774, PubMed:19759537, PubMed:21478859, PubMed:23622245, PubMed:25043379). Acts as an activator of the Wnt signaling pathway by mediating poly-ADP-ribosylation of AXIN1 and AXIN2, 2 key components of the beta-catenin destruction complex: poly-ADP-ribosylated target proteins are recognized by RNF146, which mediates their ubiquitination and subsequent degradation (PubMed:19759537, PubMed:21478859). Also mediates poly-ADP-ribosylation of BLZF1 and CASC3, followed by recruitment of RNF146 and subsequent ubiquitination (PubMed:21478859). Mediates poly-ADP-ribosylation of TERF1, thereby contributing to the regulation of telomere length (PubMed:11739745). Stimulates 26S proteasome activity (PubMed:23622245).
Cellular Location	Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Nucleus. Chromosome, telomere Note=Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4- vesicles. Also found around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS2 is found in the nucleus, associated with TRF1
Tissue Location	Highly expressed in placenta, skeletal muscle, liver, brain, kidney, heart, thymus, spinal cord, lung, peripheral blood leukocytes, pancreas, lymph nodes, spleen, prostate, testis, ovary, small intestine, colon, mammary gland, breast and breast carcinoma, and in common-type meningioma. Highly expressed in fetal liver, heart and brain

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Tankyrase 2) Polyclonal Antibody, Unconjugated (AP58658) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.