

BCCIP Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58712

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9P287
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35979

Additional Information

Gene ID	56647
Other Names	BRCA2 and CDKN1A-interacting protein, P21- and CDK-associated protein 1, Protein TOK-1, BCCIP, TOK1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	BCCIP
Synonyms	TOK1
Function	During interphase, required for microtubule organizing and anchoring activities. During mitosis, required for the organization and stabilization of the spindle pole (PubMed: 28394342). Isoform 2/alpha is particularly important for the regulation of microtubule anchoring, microtubule stability, spindle architecture and spindle orientation, compared to isoform 1/beta (PubMed: 28394342). May promote cell cycle arrest by enhancing the inhibition of CDK2 activity by CDKN1A. May be required for repair of DNA damage by homologous recombination in conjunction with BRCA2. May not be involved in non-homologous end joining (NHEJ).
Cellular Location	Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, spindle pole. Note=Colocalizes with BRCA2 in discrete nuclear foci (PubMed:15713648). In interphase, preferential localizes to the mother centriole (PubMed:28394342).

Recruited to the spindle pole matrix and centrosome by microtubules and dynein/dynactin activity (PubMed:28394342). [Isoform 2]: Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Note=Isoform 2/alpha tends to be more abundant at, and more strongly associated with, centrosomes than isoform 1/beta.

Tissue Location

Expressed at high levels in testis and skeletal muscle and at lower levels in brain, heart, kidney, liver, lung, ovary, pancreas, placenta, and spleen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.