

Importin4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58725

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q8TEX9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	118715

Additional Information

Gene ID	79711
Other Names	Importin-4, Imp4, Importin-4b, Imp4b, Ran-binding protein 4, RanBP4, IPO4, IMP4B, RANBP4
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	IPO4
Synonyms	IMP4B, RANBP4
Function	Nuclear transport receptor that mediates nuclear import of proteins, such as histones, RPS3A, TNP2 and VDR (PubMed: 11823430 , PubMed: 16207705 , PubMed: 17682055 , PubMed: 21454524 , PubMed: 30177573). Serves as receptor for nuclear localization signals (NLS) in cargo substrates (PubMed: 11823430 , PubMed: 16207705). Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism (PubMed: 11823430 , PubMed: 16207705). At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran (PubMed: 11823430). The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed: 11823430). Mediates the nuclear import of the histone H3-H4 dimer

when in complex with ASF1 (ASF1A or ASF1B) (PubMed:[21454524](#), PubMed:[29408485](#), PubMed:[36103578](#)). Also mediates the nuclear import of monomeric histones H3.1 and H4 (PubMed:[30177573](#)). Mediates the ligand-independent nuclear import of vitamin D receptor (VDR) (PubMed:[16207705](#)). In vitro, mediates the nuclear import of human cytomegalovirus UL84 by recognizing a non- classical NLS (PubMed:[12610148](#)).

Cellular Location

Cytoplasm. Nucleus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.