

UCHL5IP Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58750

Product Information

| | |
|--------------------------|-------------------------|
| Application | WB, IHC-P, IHC-F, IF, E |
| Primary Accession | Q99871 |
| Reactivity | Rat, Dog, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 39794 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 55559 |
| Other Names | HAUS augmin-like complex subunit 7, 26S proteasome-associated UCH37-interacting protein 1, UCHL5-interacting protein, X-linked protein STS1769, HAUS7, UCHL5IP, UIP1 |
| Dilution | WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000 |
| Format | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce |
| Storage | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

Protein Information

| | |
|--------------------------|---|
| Name | HAUS7 |
| Synonyms | UCHL5IP, UIP1 |
| Function | Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex. |
| Cellular Location | Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Localizes to interphase centrosomes and to mitotic spindle microtubules. |
| Tissue Location | Detected in spleen, thymus, testis, ovary, small intestine and colon, with highest levels of expression in testis and ovary. |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.