

SKA2 Rabbit pAb

SKA2 Rabbit pAb
Catalog # AP58773

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8WVK7
Predicted	Rat, Pig, Dog, Human, Mouse, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14188
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SKA2
Epitope Specificity	2-88/121
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, cytoskeleton, spindle.Chromosome, centromere, kinetochore. Note=Localizes to the outerkinetochore and spindle microtubules during mitosis in a NDC80complex-dependent manner. Localizes to both the mitotic spindle andkinetochore-associated proteins.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Ska2 (spindle and kinetochore associated complex subunit 2), also known as FAM33A, is a 121 amino acid component of the Ska1 complex, a microtubule-binding subcomplex of the outer kinetochore that is critical for proper chromosome segregation. The Ska1 complex is a component of the kinetochore-microtubule interface and directly associates with microtubules as oligomeric assemblies. Localized to the outer kinetochore and spindle microtubules during cell proliferation, Ska2 is essential for spindle checkpoint silencing and exit from mitosis. Downregulation of Ska2 leads to delayed recruitment of MAD2, a component of the mitotic spindle checkpoint, to several kinetochores resulting in occasional loss of individual chromosomes from the metaphase plate. Ska2 is encoded by a gene located on human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

Additional Information

Gene ID	348235
Other Names	Spindle and kinetochore-associated protein 2, Protein FAM33A, SKA2, FAM33A
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SKA2
Synonyms	FAM33A
Function	Component of the SKA1 complex, a microtubule-binding subcomplex of the outer kinetochore that is essential for proper chromosome segregation (PubMed:17093495, PubMed:19289083, PubMed:23085020). Required for timely anaphase onset during mitosis, when chromosomes undergo bipolar attachment on spindle microtubules leading to silencing of the spindle checkpoint (PubMed:17093495). The SKA1 complex is a direct component of the kinetochore-microtubule interface and directly associates with microtubules as oligomeric assemblies (PubMed:19289083). The complex facilitates the processive movement of microspheres along a microtubule in a depolymerization- coupled manner (PubMed:17093495, PubMed:19289083). In the complex, it is required for SKA1 localization (PubMed:19289083). Affinity for microtubules is synergistically enhanced in the presence of the ndc-80 complex and may allow the ndc-80 complex to track depolymerizing microtubules (PubMed:23085020).
Cellular Location	Cytoplasm, cytoskeleton, spindle. Chromosome, centromere, kinetochore. Note=Localizes to the outer kinetochore and spindle microtubules during mitosis in a NDC80 complex-dependent manner. Localizes to both the mitotic spindle and kinetochore- associated proteins.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.