

TBRG4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58798

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<u>Q969Z0</u>
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70738

Additional Information

Gene ID	9238
Other Names	FAST kinase domain-containing protein 4, Cell cycle progression restoration protein 2, Cell cycle progression protein 2, Protein TBRG4, Transforming growth factor beta regulator 4 {ECO:0000312 HGNC:HGNC:17443}, TBRG4 (<u>HGNC:17443</u>)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000 -10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information	
Name	TBRG4 (<u>HGNC:17443</u>)
Function	Plays a role in processing of mitochondrial RNA precursors and in stabilization of a subset of mature mitochondrial RNA species, such as MT-CO1, MT-CO2, MT-CYB, MT-CO3, MT-ND3, MT-ND5 and MT-ATP8/6. May play a role in cell cycle progression (PubMed: <u>9383053</u>).
Cellular Location	Mitochondrion matrix
Tissue Location	Ubiquitously expressed (PubMed:12759187). Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in smooth muscle (PubMed:20869947)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.