

# **CA7 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58822

### **Product Information**

**Application** WB, IHC-P, IHC-F, IF, E

Primary Accession P43166

**Reactivity** Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 29658
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human CA7

**Epitope Specificity** 86-150/264

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Cytoplasm (Probable).

**SIMILARITY** Belongs to the alpha-carbonic anhydrase family.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Carbonic anhydrases are a large family of zincmetalloenzymes that catalyze

the reversible hydration of carbondioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, boneresorption, and the formation of aqueous humor, cerebrospinalfluid, saliva, and gastric acid. They show extensive diversity intissue distribution and in their subcellular localization. Thecytosolic protein encoded by this gene is predominantly expressed in the salivary glands. Alternative splicing in the

coding regionresults in multiple transcript variants encoding

differentisoforms. [provided by RefSeq, Jul 2008].

#### **Additional Information**

Gene ID 766

Other Names Carbonic anhydrase 7, 4.2.1.1, Carbonate dehydratase VII, Carbonic

anhydrase VII, CA-VII, CA7

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000

-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


## **Protein Information**

Name CA7

**Function** Reversible hydration of carbon dioxide.

Cellular Location Cytoplasm.

## **Images**



Sample:

Raw264.7(Mouse) Cell Lysate at 30 ug Kidney (Mouse) Lysate at 40 ug

Primary: Anti- CA7 (AP58822) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at

1/20000 dilution

Predicted band size: 30 kD Observed band size: 30 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.