

ENC1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58844

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O14682
Reactivity	Mouse, Rat
Predicted	Human, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66130
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ENC1
Epitope Specificity	241-300/589
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus matrix. Cytoplasm, cytoskeleton.
SIMILARITY	Contains 1 BTB (POZ) domain. Contains 6 Kelch repeats.
SUBUNIT	Binds to RB1. Hypophosphorylated RB1 associates with ENC1 during neuronal differentiation, while hyperphosphorylated RB1 associates with ENC1 in undifferentiating cells. Part of a complex that contains CUL3, RBX1 and ENC1. Ubiquitinated and probably targeted for proteasome-independent degradation.
Post-translational modifications	
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Actin-binding protein involved in the regulation of neuronal process formation and in differentiation of neural crest cells. May be down-regulated in neuroblastoma tumors. Substrate-specific adapter of an E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Tissue specificity: Detected in fetal brain tissue, moderate expression in fetal heart, lung and kidney. Highly expressed in adult brain, particularly high in the hippocampus and amygdala, and spinal chord. Detectable in adult pancreas.

Additional Information


Gene ID	8507
Other Names	Ectoderm-neural cortex protein 1, ENC-1, Kelch-like protein 37, Nuclear matrix protein NRP/B, p53-induced gene 10 protein, ENC1, KLHL37, NRPB, PIG10
Target/Specificity	Detected in fetal brain tissue, moderate expression in fetal heart, lung and kidney. Highly expressed in adult brain, particularly high in the hippocampus and amygdala, and spinal chord. Detectable in adult pancreas.

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:50,IF=1:100-500,Flow-Cyt=0.2ug/test,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ENC1
Synonyms	KLHL37, NRPB, PIG10
Function	Actin-binding protein involved in the regulation of neuronal process formation and in differentiation of neural crest cells. Down- regulates transcription factor NF2L2/NRF2 by decreasing the rate of protein synthesis and not via a ubiquitin-mediated proteasomal degradation mechanism.
Cellular Location	Nucleus matrix. Cytoplasm. Cytoplasm, cytoskeleton
Tissue Location	Detected in fetal brain tissue, moderate expression in fetal heart, lung and kidney. Highly expressed in adult brain, particularly high in the hippocampus and amygdala, and spinal cord Detectable in adult pancreas. May be down-regulated in neuroblastoma tumors

Images



Blank control: A549.

Primary Antibody (green line): Rabbit Anti-ENC1 antibody (AP58844)

Dilution: 0.2 µg /10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG .


Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 µg /test.


Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.


Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ENC1) Polyclonal Antibody, Unconjugated (AP58844) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit)



(sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ENC1) Polyclonal Antibody, Unconjugated (AP58844) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: A549.

Primary Antibody (green line): Rabbit Anti-ENC1 antibody (AP58844)

Dilution: 1 µg /10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 3 µg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.